

Cat. No: P2091, 100 rxns/20- μ l rxn
P2092, 500 rxns/20- μ l rxn

SYBR[®] Green qPCR Mix

For research use only

Components

Component	P2091	P2092
2X SYBR [®] Green qPCR Mix	1 ml	1 ml \times 5
Nuclease-free Water	1 ml	1 ml \times 5

Storage

This reagent can be stored for 2 months at 4°C and protected from light. For longer storage, it should be kept at -20°C and protected from light.

Description

SYBR[®] Green qPCR Mix is designed for high-performance, high-throughput real-time PCR. The kit contains Taq DNA Polymerase engineered through a process of molecular evolution. The result is a unique polymerase that specifically designed for qPCR using SYBR[®] Green I chemistry dye.

2X SYBR[®] Green qPCR Mix is a convenient premix of the components (except primers and DNA template) that necessary to perform real-time polymerase chain reaction (PCR) using SYBR[®] Green I dye with enhanced sensitivity and specificity. The SYBR[®] Green I dye binds to double-stranded DNA (dsDNA), thus providing a fluorescent signal that reflects the amount of dsDNA products generated during PCR.

The reaction system of this product can be prepared at room temperature without an ice box. The prepared PCR reaction system can be placed at room temperature for 24 hours and the amplification efficiency remains unchanged.

Applications

- Gene expression analysis
- Low-copy gene detection
- Microarray validation
- Gene knockdown validation

Features

- Compatible with many Real-time systems which not require ROX reference dye
- Hot-start technology brings high specificity and reproducible amplification

Composition of the 2X SYBR Green qPCR Mix

100mM KCl , 5mM MgCl₂, 400 μ M dNTPs, 0.1U/ μ l Hotstart Taq DNA Polymerase, 1 \times SYBR[®] Green I and other optimized buffer components.

Table of Instrument Guide

Instruments	Final Conc. of ROX
ABI PRISM 7000/ PRISM 7700/ 7300/ 7900HT/ StepOne/ StepOnePlus/ GeneAmp 5700	500 nM, high ROX
ABI 7500/ 7500 Fast/ ViiA 7/ QuantStudio 6/7/12K Flex; Agilent Stratagene Mx3000P/ Mx3005P/ Mx4000	50 nM, low ROX
Bio-Rad CFX96/ CFX384/ iQ/ iQ5; MJ Research Opticon 2/ Chromo 4; Roche LightCycler 480/ 96; Corbett Rotor Gene G/ Q/ 3000/ 6000; Thermo PikoReal 96; Eppendorf MasterCycler ep realplex; Cepheid Smart Cyclor	No ROX

Protocol

Note: Please follow the procedures outlined in the manual of each respective instrument.

1. Preparation of reaction solution (Take the Biorad CFX96 as an example)

Add the following reagents to the proper thermal cycler reaction tube or plate on ice:

Component	20- μ l rxn	Final Conc.
2X SYBR [®] Green qPCR Mix	10 μ l	1X
Forward Primer (10 μ M)	0.4 μ l	0.2 μ M
Reverse Primer (10 μ M)	0.4 μ l	0.2 μ M
Template DNA	variable	0.05-5ng/ μ l
Water, nuclease-free	to 20 μ l	–

Note:

- The primer concentration can be further optimized. The optimal range for primers is 0.1~1 μ M.
- Prepare according to the recommended volume of each instrument.
- Use 1-10ng cDNA or 10-100ng gDNA for each reaction.
- Users can increase the amount of the the qPCR Mix when using low-copy gene as template.
- Users can reduce the amount of the qPCR Mix, if the melting curve comes with impure peaks.

2. Setup the plate

Transfer the reaction mixture to PCR tubes/plates. Reaction volumes can be reduced to 10 μ l if the instrument supports a low volume system.

Cap or seal the reaction tubes/plates then centrifuge briefly to spin down the contents and eliminate any air bubbles.

3. Preform qPCR using the following thermal cycling condition

Set the thermal cycling conditions using default PCR thermal cysling conditions specified in the following tables according to the instrument cycling parameters and melting temperatures of the specific primers.

2-step PCR mode:

Initial Denaturation	95°C	1 min	Holding Stage
Denaturation	95°C	5 sec	Cycling Stage 40 Cycles
Annealing & Extension	60°C ^[1]	20~60 sec ^[2]	
Melting curve analysis (optional) ^[3]			

3-step PCR mode:

Initial Denaturation	95°C	1 min	Holding Stage
Denaturation	95°C	5 sec	Cycling Stage 40 Cycles
Annealing	55°C ^[1]	30 sec	
Extension	72°C	30~60 sec ^[2]	
Melting curve analysis (optional) ^[3]			

Note:

[1] To improve the specificity, the annealing temperature can be increased. Set up signal collection in the Annealing&Extension or Extension phases respectively.

[2] Consider the instrument type when setting the extension time. If the amplification efficiency needs to be improved, the extension time can be increased.

[3] Different instruments have different melting curve acquisition procedures, generally according to the default melting curve acquisition procedures of the instrument.

4. Analyze the results

Data analysis varies depending on the instrument used. Please refer to your instrument user guide for information.

Important Notes**Template**

Genomic DNA, plasmid DNA, or cDNA can be used as template. For optimal quantitative results use up to 20ng of genomic DNA or plasmid DNA per 20 µl reaction (for smaller volumes, the amount of template should be decreased equivalently). Using greater amounts of template may reduce the maximum fluorescence signal and linearity of standard curves due to binding of the SYBR[®] Green I dye to the template. For two-step RT-PCR, use either undiluted or diluted cDNA generated from up to 1µg of total RNA. The volume of the cDNA added (from the RT reaction) should not exceed 10% of the final PCR volume (e.g., for a 20 µl qPCR reaction, use up to 2.0 µl of undiluted cDNA).

Primers

Careful primer design and purification (HPLC-purified primers are recommended) is particularly important in order to minimize loss in sensitivity due to the production of nonspecific amplification products in SYBR[®] Green I-based qPCR. This effect becomes more prominent at low target concentrations. To maximize the sensitivity of the assay, use the lowest concentration of primers that can be used without compromising the efficiency of the PCR reaction (50-400nM of each primer). For optimal results, design primers that amplify PCR products 60-400 bp in length. The primers should exhibit a melting temperature (T_m) of approximately 60°C. We recommend designing primers specifically for amplification of cDNA derived from mRNA. This prevents

amplification of contaminating genomic DNA and inaccurate quantification of mRNA.

Melting Curve Analysis

Following real-time qPCR, melting curve analysis should always be performed to identify the presence of primer-dimers and analyze the specificity of the reaction. Program your thermocycler according to the instructions provided.

Quality Control

The absence of endodeoxyribonucleases, exodeoxyribonucleases and ribonucleases is confirmed by appropriate quality tests. Functionally tested in amplification of a single-copy gene from human genomic DNA.

Product Use Limitations

SYBR[®] Green qPCR Mix is sold exclusively for research purposes and in vitro use. Neither the product, nor any individual components, was tested for use in diagnostic applications or for drug development, nor is it suitable for administration to humans or animals. Please refer to the MSDS, available upon request.