

Xho I

Instruction for Use

Cat. No./Spec.: E1029-A/400 rxns, E1029-B/1,200 rxns

Recognition site:

5' C ↓ T C G A G 3'
3' G A G C T ↑ C 5'

Product Description

The Xho I restriction endonuclease recognizes the C[^]TCGAG site and achieves optimal cutting efficiency within 5 – 15 minutes at 37 ° C using the universal buffer. Isoschizomers: PaeR7 I , Sfr274 I , Sla I , Str I , Tli I .

GDSBio restriction endonucleases exhibit 100% activity in both Digest and Green reaction buffers.

The universal Digest buffer allows for rapid single-enzyme, double-enzyme, or multiple-enzyme digestion of DNA within 5–15 minutes, eliminating the need for buffer exchange or subsequent DNA purification steps. DNA-modifying enzymes (such as the Klenow fragment, T4 DNA ligase, calf intestinal alkaline phosphatase, and T4 DNA polymerase) maintain 100% activity in the buffer. Consequently, enzymes used in downstream applications can be directly added to the reaction mixture. Shorter incubation times and the superior composition of the universal Digest buffer eliminate the star activity effect.

The Green buffer includes one density reagent and two tracking dyes for direct loading of the digestion products onto a gel.

Components

Component	E1029-A	E1029-B
Xho I	400 µL	1.2 mL
10X Digest Buffer	1 mL	1 mL × 3
10X Green Buffer	1 mL	1 mL × 3

Storage Condition & Shelf Life

Store at -20°C.

Features

- All GDSBio restriction endonucleases exhibit 100% activity in the universal buffer.
- 100% buffer compatibility with downstream applications.
- Enzymatic digestion can be completed within 5 – 15 minutes.
- Direct loading onto a gel.
- No star activity.

Scope of Application

- Molecular cloning
- Restriction site mapping
- Genotyping
- Southern blotting
- Restriction fragment length polymorphism (RFLP)
- SNP (Single Nucleotide Polymorphism) analysis

Recommended Reaction Conditions

- 1X Digest Buffer or 1X Green Buffer
- Incubate at 37°C.
- Maximum amounts for enzymatic digestion with 1 µL Xho I :
 - 1 µg lambda DNA in 5 minutes
 - 1 µg plasmid DNA in 5 minutes
 - 0.2 µg PCR product in 5 minutes
 - 1 µg genomic DNA in 10 minutes, or 5 µg genomic DNA in 60 minutes

Inactivation

Incubation at 80°C for 5 min.

Methylation Effects on Digestion

Dam: never overlaps – no effect.

Dcm: never overlaps – no effect.

CpG: completely overlaps – cleavage impaired.

EcoKI: never overlaps – no effect.

EcoBI: never overlaps – no effect.

Number of Recognition Sites in DNA

λ DNA	Φ X174	pBR322	pUC57	pUC18/19	pTZ19R/U	M13mp18/19
1	1	0	0	0	0	0

Protocol

Fast Digestion of Different DNA

① Prepare the reaction system at room temperature in the following order:

Component	Plasmid DNA	PCR product	Genomic DNA
Nuclease-free Water	15 μ L	17 μ L	30 μ L
10X Digest Buffer	2 μ L	2 μ L	5 μ L
or 10X Green Buffer			
DNA	2 μ L (up to 1 μ g)	10 μ L (~0.2 μ g)	10 μ L (5 μ g)
enzyme	1 μ L	1 μ L	5 μ L
Total volume	20 μ L	30 μ L	50 μ L

② Gently mix and briefly centrifuge.

③ Incubate at 37°C for 5 minutes (PCR products and plasmid DNA), or for 10 minutes (genomic DNA).

Optional: Heat at 80°C for 5 minutes to inactivate the enzyme.

Double and Multiple Digestion of DNA

- The total volume of enzyme in the reaction mixture should not exceed 1/10 of the total reaction volume.

- Use 1 μ L of each enzyme and scale up the reaction conditions accordingly.

- If the enzymes require different reaction temperatures, start with the enzyme that requires the lower temperature, then add the second enzyme and incubate at the higher

temperature.

Scaling up Plasmid DNA Digestion Reaction

Component	20- μ L rxn	20- μ L rxn	30- μ L rxn	40- μ L rxn	50- μ L rxn
DNA	1 μ g	2 μ g	3 μ g	4 μ g	5 μ g
enzyme	1 μ L	2 μ L	3 μ L	4 μ L	5 μ L
10X Digest Buffer or 10X Green Buffer	2 μ L	2 μ L	3 μ L	4 μ L	5 μ L
Total volume	20 μ L	20 μ L	30 μ L	40 μ L	50 μ L

Note: If the total reaction volume exceeds 20 μ L, increase the incubation time by 3-5 minutes. Use a water bath for temperature control; an air incubator is not recommended as heat transfer to the reaction mixture is slow.

This product is for research use only.